

Screening of Variables Affecting Extraction of Pectin from Unripe Banana (*Musa acuminata* × *balbisiana* var. Cardaba) Peel using Plackett–Burman Design

Receil Ann B. Tanaid*

ABSTRACT

Pectin is an important ingredient used as thickener, water binder, and stabilizer in food. Its production could be economically attractive especially if extracted from food processing wastes and by-products. This study screened the different experimental conditions that led to the significant extraction of quality pectin from Cardaba banana. The large production of banana products, which results in discarded banana peels, presents an opportunity for creating alternative processes involving this waste. The Plackett–Burman screening method was used to identify the most important factors early in the experimentation phase when complete knowledge about the factors affecting the process or method were unavailable. Seven selected variables, namely: peel nature, peel size, peel-solvent ratio, pH, extraction temperature, extraction time, and extractant:ethanol ratio, were subjected to the screening experiment. These variables are relevant to the extraction of pectin in unripe banana (*Musa acuminata* × *balbisiana* var. Cardaba) peel. Variable screening results showed that extraction time, extraction temperature, and peel:solvent ratios had significant effects on the extraction process. These factors could be further optimized to extract the best quality pectin from unripe banana peel.

Keywords: banana, Cardaba, pectin, Plackett-Burman

INTRODUCTION

Cardaba banana (*Musa acuminata* × *balbisiana* var. Cardaba) is a triploid hybrid (ABB) banana cultivar originating from the Philippines (Dela Cruz et al 2007). It is one of the most important banana varieties in Philippine cuisine, but its biggest potential is for processing into chips (Dela Cruz et al 2008) wherein unripe bananas are used. The edible portion of Cardaba banana is only 60% of the entire fruit, while the rest, specifically the banana peel, is

*Cebu Technological University, Cor. M.J. Cuenco, R. Palma Street, Cebu City, Cebu, Philippines

*Corresponding Author : Address: Department of Food Science and Technology, Cebu Technological University, Cor. M.J. Cuenco, R. Palma Street, Cebu City, Cebu, Philippines. E-mail: rtanaid19@gmail.com

Screening of Variables Affecting Extraction of Pectin from Unripe Banana Peel

considered waste (Dela Cruz 2008). However, fruit and vegetable wastes are rich sources of carbohydrates, proteins, fats, minerals, and fibers (Gowe 2015). Therefore, they have potential uses in the food industry, one of which is as food additives such as antioxidants, antimicrobials, colorants, flavorings, and thickener agents (Ayala-Zavala & González-Aguilar, 2011).

Pectin is the methylated ester of polygalacturonic acid which contains 1, 4-linked α -D-galacturonic acid residues. Pectin is capable of forming gels with sugar and acid. Because of this gelling ability, one of the well-known uses of pectin is in high-sugar jams and confectionery jellies, dating back to at least the 18th century (IPPA 2001). Because it is a natural additive for foods, pectin is being considered for a number of applications beyond the traditional jams and jellies. Pectins are now used as thickeners, water binders, and stabilizers. It is used in yogurts and pastry glazes and as a stabilizer in drinkable yogurts and blends of milk and fruit juices (May 1990). Pectin has also been investigated for its usefulness in the pharmaceutical industry, particularly stabilizing liquid pharmaceutical emulsions and suspensions.

To date, commercial pectin used for the preparation of different confections and gelling products (such as marmalades, jams, preserves, and low calorie gels) are principally produced from two pectin-rich sources, lime and apple pomace. The yield of pectin usually depends on the extraction conditions such as temperature, extraction time, pH, type of extraction solvents (Yeoh et al 2008), and many others.

The aim of this study was to screen the different experimental conditions that lead to the significant extraction of quality pectin from Cardaba banana peels. The results of the study will be subsequently used for further research on extraction optimization. As many factors can influence the pectin extraction yield and quality, Plackett–Burman design was applied to screen important variables affecting extraction of pectin from unripe banana.

MATERIALS AND METHODS

Extraction Process of Pectin from Banana Peel

Two forms of banana peel were used for pectin extraction: dried and fresh. Dried banana peel was prepared using Castillo-Israel's (2015) method, wherein fresh banana peel was dried in a forced draft oven at 55°C for 24 hours. On the other hand, fresh banana peel was thinly sliced and soaked in water prior to pectin extraction.

The extraction of pectin followed different methods based on the levels and conditions used. Banana peel was added to distilled water, pH 7 (Mohdet al 2012), and 0.50 N citric acid, pH 2 (Virk & Sogi 2004). Peel to solvent (extractant) ratio was set at 1:10 and 1:50. These were heated while

continuously stirring at different extracting temperatures (60°C and 100°C) on a hot plate for 30 and 180 minutes (Koffi et al 2013). The solution was cooled and filtered through an ordinary screen with 1-mm mesh size and two-layer cheesecloth. The filtrate was collected then added with absolute ethanol at 1:1 and 1:3 filtrate:ethanol ratio, then kept at room temperature overnight. The precipitate (ethanol-insoluble fraction) was filtered through Whatman filter paper No. 4. It was washed with 75% ethanol, then with 80% ethanol to remove soluble impurities (Azad et al 2014). The residue was oven dried for 12 hours at 55°C, cooled, and weighed.

Experimental Design for Screening Experiment

The screening experiment used the Plackett-Burman design to identify the most important factors that affected pectin extraction. Each parameter studied was explored at two (2) levels: high and low values (Table 1). Seven parameters were investigated in the experiment (Table 2) having eight (8) runs and producing different product responses, which were characterized chemically.

Table 1. High and low values for screening experiment for banana peel pectin extraction process

Variables	Peel Nature	Peel Form	Peel: Solvent Ratio	pH	Extraction Temperature (°c)	Extraction Time (Min)	Ethanol: Filtrate Ratio
+ High level	Fresh	Medium slices	1:50	7	100	180 min	1:3
- Low level	Dried	Crushed form	1:10	2	60	30 min	1:1

Table 2. Plackett-Burman 8-run design with 7-variables that would affect the extraction of pectin from Cardaba banana peel

	Peel Nature	Peel Form	Peel: Solvent Ratio	pH	Extraction Temperature (°c)	Extraction Time (min)	Ethanol: Filtrate Ratio
1	Fresh	Sliced	1:50	2	100	30	1:1
2	Fresh	Sliced	1:10	7	60	30	1:3
3	Fresh	Crushed	1:50	2	60	180	1:3
4	Dried	Sliced	1:10	2	100	180	1:3
5	Fresh	Crushed	1:10	7	100	180	1:1
6	Dried	Crushed	1:50	7	100	30	1:3
7	Dried	Sliced	1:50	7	60	180	1:1
8	Dried	Crushed	1:10	2	60	30	1:1

Screening of Variables Affecting Extraction of Pectin from Unripe Banana Peel

Characterization of Pectin

Pectin Yield

Pectin yield was calculated using the formula:

$$\text{Pectin (\%)} = \frac{\text{Weight (g) of pectin}}{\text{Weight (g) of peel taken for extraction}} \times 100$$

Moisture Content Determination

One gram of pectin sample was weighed, ground, and placed in an aluminum foil. The sample was dried in an oven for 5 hours at 100°C, cooled in a desiccator, and then weighed. This sample was not used for subsequent measurement, as pectin tends to degrade after exposure to the atmosphere. One percent was added to the percent moisture observed to conform to the Fischer method (Johnson 1945 as cited by Castillo-Israel 2015). The moisture content was determined using the formula:

$$\text{Moisture content (\%)} = \frac{\text{Weight (g) of residue}}{\text{Weight (g) of sample}} \times 100$$

Ash Content Determination

One to two (1-2) grams of pectin was ground to pass an 80-mesh screen. It was then placed into tared crucibles and was burnt in a furnace for 3-4 hours at 600°C. The ash content was determined using the formula:

$$\text{Ash content (\%)} = \frac{\text{Weight (g) of ash}}{\text{Weight (g) of sample}} \times 100$$

Determination of Equivalent Weight

The equivalent weight was determined by Ranganna's method (1995). About 0.5 g sample was placed in a 250 mL conical flask and added with 5 mL ethanol, 1g of NaCl, and 100 mL of distilled water. Finally, about six (6) drops of phenol red was added and titrated against standardized 0.1 N NaOH.

The end point was indicated by a purple or faint pink color. This neutralized solution was stored for determination of methoxyl content. The equivalent weight was calculated using the formula:

$$\text{Equivalent weight} = \frac{\text{Weight (g) of sample} \times 1000}{\text{Volume of alkali} \times \text{Normality of alkali}}$$

Determination of Methoxyl Content (MeO)

Determination of MeO was done using Ranganna's method (1995). The neutral solution was collected from determination of equivalent weight and 25 mL of 0.25 N NaOH was added. The mixed solution was stirred thoroughly and kept at room temperature for 30 minutes. After 30 minutes, 25 mL of 0.25 N HCl was added and the mixture was titrated against standardized 0.1 N NaOH. Methoxyl content was calculated using the formula:

$$\text{Methoxyl content (\%)} = \frac{\text{Volume of alkali (mL)} \times \text{Normality of alkali} \times 3.1}{\text{Weight (g) of sample}}$$

Determination of Total Anhydrouronic Acid Content (AUA)

Total AUA of pectin was calculated using the following formula (Mohamed & Hasan 1995):

$$\% \text{ of AUA} = \frac{176 \times 0.1z \times 100}{w \times 1000} + \frac{176 \times 0.1y \times 100}{w \times 1000}$$

when molecular unit of AUA (1 unit)=176 g

where,

z= mL (titre) of NaOH from Equivalent weight determination

y= mL (titre) of NaOH from methoxyl content determination

w= weight of sample

Determination of Degree of Esterification (DE)

The DE of pectin was measured on the basis of methoxyl and AUA content (Owens et al 1952) and calculated using this formula:

$$\% \text{ DE} = \frac{176 \times \% \text{MeO}}{31 \times \% \text{AUA}} \times 100$$

Statistical Analysis and Modelling

The results of physico-chemical properties analysis from the Plackett-Burman design were analyzed for variables which have significant effects using Statistica 8.0 software and Statistical Analytical Software version 9 (SAS 2008).

RESULTS AND DISCUSSION

Results of Variable Screening

Plackett-Burman was used as the experimental design to screen the important variables affecting pectin extraction from unripe banana peel. Results showed that the variables having more significant results were the peel nature, peel-solvent ratio, extraction temperature, and extraction time. The variables with less significant results were the peel size, pH, and ethanol:pectin extract ratio (Tables 3 & 4).

Controlling the moisture content of the peels was unnecessary since these were stored in an uncontrolled environment, thus contact with the surrounding's relative humidity must have affected the moisture content. The degree of moisture absorption depends on the type of dried product and the ambient conditions such as temperature, humidity, and contact time (Dauthy 1995). This was also held constant to produce a constant yield percentage in both dry and wet banana peel. The moisture content of peel used was 87.5% wet basis and 7.01% dry basis.

Table 3. Summary of effects for yield, moisture content, ash content, and equivalent weight

Parameter	Parameters			
	Yield (%)	Moisture Content (%)	Ash (%)	Equivalent Wt.
Mean/Interc.	1.26 ^{ns}	12.64 ^{ns}	13.23 ^{ns}	864.42 ^{ns}
Peel Nature	-0.22**	-0.16 ^{ns}	4.06 ^{ns}	40.84**
Peel Size	-0.11**	-0.12 ^{ns}	-1.7 ^{ns}	-55.33 ^{ns}
PS Ratio	-0.12**	-0.34*	-1.93 ^{ns}	-39.21**
pH	-0.48**	0.13 ^{ns}	0.23 ^{ns}	-69.13**
E. temp	0.89**	0.26*	-2.53 ^{ns}	144.13**
E. time	0.10**	-0.48**	-0.45 ^{ns}	-25.95**
Ethanol	-0.0 ^{ns}	-0.93**	1.04 ^{ns}	-1.97 ^{ns}

^{ns}not significant (p>0.05) *significant (p≤ 0.05) **significant (p≤ 0.01)

Table 4. Summary of effects for methoxyl content (MeO), anhydrouronic acid content (AUA), and degree of esterification (DE)

Parameter	Parameters		
	MeO (%)	AUA (%)	DE (%)
Mean/Interc.	8.16 ^{ns}	68.58 ^{ns}	67.48 ^{ns}
Peel Nature	0.35**	0.93**	1.75**
Peel Size	0.74**	5.40**	0.82**
PS Ratio	0.29**	2.64**	-0.10 ^{ns}
pH	-0.05 ^{ns}	1.23**	-1.74**
E. temp	0.55**	-0.18 ^{ns}	4.72**
E. time	0.27**	2.11**	0.10 ^{ns}
Ethanol	1.12**	6.63**	2.70**

^{ns}not significant (p>0.05) *significant (p≤ 0.05) **significant (p≤ 0.01)

For the pH variable, 0.5N citric acid with pH 2 was used, since pH 2 was found to give the highest yield of extracted pectin from table banana (Girma & Worku 2016). According to Shaha et al (2013), citric acid is the best agent for the extraction of pectin. This is in agreement with the results reported by Virk and Sogi (2004) who compared the yields of pectin extracted from apple with different acids, namely: hydrochloric acid, nitric acid, and citric acid. It was also reported by Yapo (2009) that citric acid was the least pectin degrading (depolymerizing and deesterifying) extracting agent. Therefore, it leads to pectin isolates with the best gelling properties. However, among the acids used in the experiment, it was observed that there was no significant difference in pectin yield.

Variables such as peel size and pH had negative effects on all parameters. Even though a low pH is necessary to improve the yield, the use of a strong acid solution such as hydrochloric acid could lead to smaller pectin particles owing to partial hydrolysis. Consequently, pectin solubility would increase to the point that no precipitate was formed by the addition of alcohol (Canteri-Schemin et al 2005). The peel size affects the extraction process since the crushed form (low level) contributes to the viscosity of the solution causing inefficient filtration, thus reducing the yield and compromising pectin quality. Using medium-sized peels (high level) also reduced the surface area, thus reducing the penetration rate of the solvent. For ethanol:pectin ratio, the effect on the parameters was positive, thus the high level 1:3 ratio was used.

From these results, the variables considered for possible optimization of pectin extraction were peel-solvent ratio, extraction time, and extraction temperature. Majority of the physico-chemical properties showed a positive effect to extraction temperature and time.

If the effect is positive, the closest value, either lower or higher than the indicated high level (+) concentration, is required during further optimization studies. On the other hand, if the effect is negative, the closest value, either lower or higher than the indicated low level (-) concentration, is used. This applies to the peel:solvent ratio which negatively affected majority of the parameters. These three factors were important in the extraction process since the rate of vaporization during extraction was affected by the amount of solvent, extraction temperature, and time. As the extraction process progressed, the concentration of the pectin in the solution increased causing the extraction rate to progressively decrease.

Increase in viscosity of the solution (due to vaporization) led to the reduction of pectin extract, resulting in a lower yield when precipitated with alcohol. High temperature and longer boiling time encourage energy and water (or solvent) loss through vaporization (Girma & Worku 2016). The peel:solvent ratio must be optimized to attain a pectin extract that, if precipitated with alcohol, will produce the highest yield possible.

CONCLUSION AND RECOMMENDATIONS

The Plackett-Burman variable screening was used to identify the variables that would significantly affect pectin extraction. A 7-variable, 8-run screening experiment was conducted with the following variables: peel nature, peel size, peel:solvent ratio, pH, extraction temperature, extraction time, and extractant:ethanol ratio. Values for extraction temperature and time showed significant positive effect, whereas peel:solvent ratio showed significant negative effect on majority of the physico-chemical tests. The results identified extraction time, extraction temperature, and peel:solvent ratio as important factors in pectin extraction. However, these have to be optimized to produce pectin of the best quality. Optimization of extraction temperature, extraction time, and peel:solvent ratio is suggested to produce the best quality of pectin from Cardaba banana peel.

REFERENCES

- Azad AKM, Ali MA, Sorifa Akter MST, Rahman J & Ahmed M. 2014. Isolation and Characterization of Pectin Extracted from Lemon Pomace during Ripening. *Journal of Food and Nutrition Sciences*. Vol. 2, No. 2, pp. 30-35. doi: 10.11648/j.jfns.20140202.12. Retrieved from: <http://www.sciencepublishinggroup.com/j/jfns> on July 13, 2016
- Ayala-Zavala JF & González-Aguilar, G. A. 2011. Use of additives to preserve the quality of fresh-cut fruits and vegetables. *Advances in fresh-cut fruits and vegetables processing*. Boca Raton, FL, USA: CRC Press. pp. 231-254
- Braddock, R.J. 1999. *Handbook of citrus by-products and processing technology*. John Wiley & Sons, Inc. New York. 191-7.
- Castillo-Israel KAT, Bagueo SF, Diasanta MDB, Lizardo RCM, Dizon EL & Mejico MIF. 2015. Extraction and characterization of pectin from Saba banana [*Musa 'saba' (Musa acuminata x Musa balbisiana)*] peel wastes: A preliminary study. *International Food Research Journal* 22(1): 202-207. University of the Philippines Los Baños, Laguna, Philippines
- Canteri-Schemin MH, Fertoni HCR, Waszczyński N & Wosiacki G. 2005. Extraction of pectin from apple pomace, *Brazilian Archives of Biology and Technology*, 48, 259-266
- Dauthy ME. 1995. *Fruit and vegetable processing*. FAO Agricultural Services Bulletin. Cited: May 27, 2017. Retrieved from: www.fao.org/docrep/V5030E/V5030E0b.htm
- Dela Cruz FS, Gueco LS, Damasco OP, Banasihan IG, Lladones RV, Van Den Bergh I, & Molina AB. 2007. *Catalogue of Introduced and Local Banana Cultivars in the Philippines*. Biodiversity International, and DABAR. University of the Philippines Los Baños. 63 pp.
- Girma E and Worku T. 2016. Extraction and Characterization of Pectin from

- Selected Fruit Peel Waste. International Journal of Scientific and Research Publications, 6 (2): 447-449
- Gowe C. 2015. Review on Potential Use of Fruit and Vegetables By-Products as A Valuable Source of Natural Food Additives. Food Science and Quality Management. 45:47-61. Cited: July 13, 2016. Availbale at: www.iiste.org
- IPPA: International Pectin Producers Association. 2001. What is pectin? Retrieved at: http://www.ippa.info/commercial_production_of_pectin.htm. On July 13, 2016
- Koffi KL, Yapo BM & Besson V. 2013. Extraction and characterization of gelling pectin from the peel of *Poncirus trifoliata* fruit. Agricultural Sciences. Vol. 4, No. 11, 614-619. Retrieved from: <http://dx.doi.org/10.4236/as.2013.411082> on July 13, 2016
- Long-Shan TL, Chieh-Chang P & Bo-Kun T. 2003. The influence of medium design on lovastatin production and pellet formation with a high-producing mutant of *Aspergillus terreus* in submerged cultures. Process Biochemistry 38: 1317–1326.
- May CD. 1990. Industrial Pectins: Sources, Production and Applications, Carbohydrate Polymers, 12:79-99.
- Mohd NS, Ramli N, Mohd N & Meon Z. 2012. Extraction and Characterization of Pectin from Dragon Fruit (*Hylocereus polyrhizus*) using Various Extraction Conditions. Sains Malaysiana 41(1): 41–45
- Owens HS, Mccready RM, Shepherd AD, Schultz SH, Phippen EL, Swenson HA, Miers JC, Erlandsen RF & Maclay WD. 1952. Methods used at Western Regional Research Laboratory for extraction and analysis of pectic materials. AIC-340, Western Regional Research Laboratory, Albany, California.
- Srinivas MRS, Chand N & Lonsane BK. 1994. Use of Plackett–Burman design for rapid screening of several nitrogen sources, growth/product promoters, minerals and enzyme inducers for the production of alpha-galactosidase by *Aspergillus niger* MRSS 234 in solid state fermentation. Bioprocess. Eng. 10:139–144.
- Virk BS and Sogi DS. 2004. Extraction and characterization of pectin from apple pomace (*Malus pumila* cv. Amri) peel waste, International Journal of Food Properties, 7:1-11
- Yapo BM, Robert C, Etienne I, Wathelet B & Paquot M. 2007. Effect of extractions on the yield, purity and surface properties of sugar beet pulp pectin extracts. *Food chemistry*, 100:1356-1364.
- Yeoh S, Shi J & Langrish TAG. 2008. Comparisons between different techniques for water-based extraction of pectin from orange peels. Desalination 218: 229-237.